

EP 04 11008



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

REC'D 18 OCT 2004

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03023288.8

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 03023288.8
Demande no:

Anmeldetag:
Date of filing: 15.10.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Bayer HealthCare AG
Kaiser-Wilhelm-Allee
51373 Leverkusen
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Urea derivatives

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C07C275/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

EPO - Mur.
41
15. Okt. 200**UREA DERIVATIVES****DETAILED DESCRIPTION OF INVENTION****TECHNICAL FIELD**

The present invention relates to an urea derivative which is useful as an active ingredient of
5 pharmaceutical preparations. The urea derivative of the present invention has vanilloid receptor
(VR1) antagonistic activity, and can be used for the prophylaxis and treatment of diseases
associated with VR1 activity, in particular for the treatment of overactive bladder, urinary
incontinence such as urge urinary incontinence, chronic pain, neuropathic pain, postoperative pain,
rheumatoid arthritic pain, neuralgia, neuropathies, algesia, nerve injury, ischaemia, neuro-
10 degeneration, stroke, and inflammatory disorders such as asthma, and chronic obstructive
pulmonary (or airways) disease (COPD).

BACKGROUND ART

Vanilloid compounds are characterized by the presence of vanillyl group or a functionally
equivalent group. Examples of several vanilloid compounds or vanilloid receptor modulators are
15 vanillin (4-hydroxy-3-methoxy-benzaldehyde), guaiacol (2-methoxy-phenol), zingerone (4-/4-
hydroxy-3-methoxyphenyl/-2-butanon), eugenol(2-methoxy4-/2-propenyl/phenol), and capsaicin
(8-methy-N-vanillyl-6-noneneamide).

Among others, capsaicin, the main pungent ingredient in "hot" chili peppers, is a specific
neurotoxin that desensitizes C-fiber afferent neurons. Capsaicin interacts with vanilloid receptors
20 (VR1), which are predominantly expressed in cell bodies of dorsal root ganglia (DRG) or nerve
endings of afferent sensory fibers including C-fiber nerve endings [Tominaga M, Caterina MJ,
Malmberg AB, Rosen TA, Gilbert H, Skinner K, Raumann BE, Basbaum AI, Julius D: The cloned
capsaicin receptor integrates multiple pain-producing stimuli. Neuron. 21: 531-543, 1998]. The
VR1 receptor was recently cloned [Caterina MJ, Schumacher MA, Tominaga M, Rosen TA,
25 Levine JD, Julius D: Nature 389: 816-824, (1997)] and identified as a nonselective cation channel
with six transmembrane domains that is structurally related to the TRP (transient receptor
potential) channel family. Binding of capsaicin to VR1 allows sodium, calcium and possibly
potassium ions to flow down their concentration gradients, causing initial depolarization and
release of neurotransmitters from the nerve terminals. VR1 can therefore be viewed as a
30 molecular-integrator of chemical and physical stimuli that elicit neuronal signals in pathological
conditions or diseases.

There is abundant direct or indirect evidence that shows the relation between VR1 activity and diseases such as pain, ischaemia, and inflammatory disorders (e.g., WO 99/00115 and 00/50387). Further, it has been demonstrated that VR1 transduces reflex signals that are involved in the overactive bladder of patients who have damaged or abnormal spinal reflex pathways [De Groat WC: A neurologic basis for the overactive bladder. Urology 50 (6A Suppl): 36-52, 1997].
5 Desensitisation of the afferent nerves by depleting neurotransmitters using VR1 agonists such as capsaicin has been shown to give promising results in the treatment of bladder dysfunction associated with spinal cord injury and multiple sclerosis [(Maggi CA: Therapeutic potential of capsaicin-like molecules - Studies in animals and humans. Life Sciences 51: 1777-1781, 1992) and
10 (DeRidder D; Chandiramani V; Dasgupta P; VanPoppel H; Baert L; Fowler CJ: Intravesical capsaicin as a treatment for refractory detrusor hyperreflexia: A dual center study with long-term followup. J. Urol. 158: 2087-2092, 1997)].

It is anticipated that antagonism of the VR1 receptor would lead to the blockage of neurotransmitter release, resulting in prophylaxis and treatment of the conditions and diseases
15 associated with VR1 activity.


It is therefore expected that antagonists of the VR1 receptor can be used for prophylaxis and treatment of the conditions and diseases including chronic pain, neuropathic pain, postoperative pain, rheumatoid arthritic pain, neuralgia, neuropathies, algnesia, nerve injury, ischaemia, neurodegeneration, stroke, inflammatory disorders, urinary incontinence (UI) such as urge urinary
20 incontinence (UUI), and/or overactive bladder.

UI is the involuntary loss of urine. UUI is one of the most common types of UI together with stress urinary incontinence (SUI) which is usually caused by a defect in the urethral closure mechanism. UUI is often associated with neurological disorders or diseases causing neuronal damages such as dementia, Parkinson's disease, multiple sclerosis, stroke and diabetes, although it also occurs in
25 individuals with no such disorders. One of the usual causes of UUI is overactive bladder (OAB) which is a medical condition referring to the symptoms of frequency and urgency derived from abnormal contractions and instability of the detrusor muscle.

There are several medications for urinary incontinence on the market today mainly to help treating UUI. Therapy for OAB is focused on drugs that affect peripheral neural control mechanisms or
30 those that act directly on bladder detrusor smooth muscle contraction, with a major emphasis on development of anticholinergic agents. These agents can inhibit the parasympathetic nerves which control bladder voiding or can exert a direct spasmolytic effect on the detrusor muscle of the bladder. This results in a decrease in intravesicular pressure, an increase in capacity and a

reduction in the frequency of bladder contraction. Orally active anticholinergic drugs such as propantheline (ProBanthine), tolterodine tartrate (Detrol) and oxybutynin (Ditropan) are the most commonly prescribed drugs. However, their most serious drawbacks are unacceptable side effects such as dry mouth, abnormal visions, constipation, and central nervous system disturbances. These side effects lead to poor compliance. Dry mouth symptoms alone are responsible for a 70% non compliance rate with oxybutynin. The inadequacies of present therapies highlight the need for novel, efficacious, safe, orally available drugs that have fewer side effects.

WO03/014064 discloses the compounds represented by the general formula:



10 wherein

X represents C₃₋₈ cycloalkyl optionally fused by benzene, optionally substituted naphthyl, optionally substituted phenyl, optionally substituted phenyl C₁₋₆ straight alkyl, phenyl fused by cycloalkyl, etc;

Q^{aa} represents CH or N;


15 R^{aa} represents hydrogen or methyl;

R^{bb} represents hydrogen or methyl; and

Y represents substituted naphthyl,

as a vanilloid receptor antagonist.

WO03/022809 discloses the compounds having vanilloid receptor antagonist activity represented by the general formula:




wherein

P and P' independently represent aryl or heteroaryl;

R^{a1} and R^{a2} independently represent hydrogen, alkoxy, hydroxy, etc;

5 n is 0, 1, 2 or 3; p and q are independently 0, 1, 2, 3 or 4; r is 1, 2 or 3; and s is 0, 1 or 2.

WO03/053945 discloses the compounds having vanilloid receptor antagonist activity represented by the general formula:




wherein

10 P^a represents phenyl, naphthyl or heterocyclyl;

n is 2, 3, 4, 5 or 6; p is independently 0, 1, 2, 3 or 4;


R^{b1} represents hydrogen, alkoxy, hydroxy, etc; and

R^{a2} represents



15 wherein X is a bond, C, O, or NR^{b8}; and r, q, R^{b3}, R^{b4} are defined in the application.

WO03/070247 discloses the compounds having vanilloid receptor antagonist activity represented by the general formula:




wherein

- 5 Xc_1 represents N or CR^{c1} ; Xc_2 represents N or CR^{c2} ; Xc_3 represents N, NR^{c3} or CR^{c3} ; Xc_4 represents a bond, N or CR^{c4} ; Xc_5 represents N or C; provided that at least one of Xc_1 , Xc_2 , Xc_3 and Xc_4 is N; Zc_1 represents O, NH or S; Zc_2 represents a bond, NH or S; L^c represents alkylene, cycloalkylene, etc; R^{c1} , R^{c2} , R^{c3} , R^{c4} , R^{c5} , R^{c6} , R^{c7} , R^{c8a} , R^{c8b} are defined in the application; and R^{c9} represents hydrogen, aryl, cycloalkyl, and heterocycle.
- 10 However, the development of a compound which has effective VR1 antagonistic activity and can be used for the prophylaxis and treatment of diseases associated with VR1 activity, in particular for the treatment of urinary incontinence, urge urinary incontinence, overactive bladder as well as pain, and/or inflammatory diseases such as asthma and COPD has been desired.

SUMMARY OF THE INVENTION

- 15 This invention is to provide an urea derivative of the formula (I), their tautomeric and stereoisomeric form, and salts thereof:



wherein

n represents an integer of 0 to 6;

20 Q_1 and Q_4 independently represent direct bond or methylene;

Chemical bond between Q_2 --- Q_3 is selected from the group consisting of a single bond and a double bond;

when Q_2 --- Q_3 is a single bond, Q_2 represents CHR^2 , or CO, and Q_3 represents CHR^3 ,

when Q_2 --- Q_3 is a double bond, Q_2 represents CR^2 and Q_3 represents CR^3 ;

5 wherein

R^2 represents hydrogen, hydroxy, C_{1-6} alkoxy or C_{1-6} alkanoyloxy;

R^3 represents hydrogen, hydroxy, C_{1-6} alkoxy, C_{1-6} alkanoyloxy, or C_{1-6} alkyl optionally substituted by hydroxy, C_{1-6} alkoxy or C_{1-6} alkanoyloxy,

with the proviso that Q_1 and Q_4 can not be direct bond at the same time;

10 R^2 and R^3 can not be hydrogen at the same time;

when Q_1 and Q_4 are both methylene and R^3 is hydroxy, R^2 is hydroxy, C_{1-6} alkoxy or C_{1-6} alkanoyloxy;

when Q_1 is direct bond, R^2 is hydroxy, C_{1-6} alkoxy or C_{1-6} alkanoyloxy; and when Q_4 is direct bond, R^2 is hydrogen, C_{1-6} alkoxy or C_{1-6} alkanoyloxy;

15 and

R^4 represents aryl optionally having one or two substituents selected from the group consisting of halogen, hydroxy, C_{1-6} alkylamino, di(C_{1-6} alkyl)amino, C_{3-8} cycloalkylamino, C_{1-6} alkoxy carbonyl, phenyl, benzyl, sulfonamide, C_{1-6} alkanoyl, C_{1-6} alkanoylamino, carbamoyl, C_{1-6} alkylcarbamoyl, cyano, C_{1-6} alkyl optionally substituted by cyano, C_{1-6} alkoxy carbonyl, or mono-, di-, or tri-halogen, C_{1-6} alkoxy optionally substituted by mono-, di-, or tri- halogen, phenoxy optionally substituted by halogen or C_{1-6} alkyl, and C_{1-6} alkylthio optionally substituted by mono-, di-, or tri- halogen.

20

In another embodiment, the urea derivatives of formula (I) are those wherein;

25 Q_1 and Q_4 represent methylene;

Q_2 --- Q_3 is a single bond;

Q₂ represents CHR², or CO,

wherein

R² represents hydroxy, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy; and

Q₃ represents CHR³,

5 wherein

R³ represents hydrogen, hydroxy, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy.

In another embodiment, the urea derivatives of formula (I) are those wherein;

Q₁ represents methylene;

Q₄ represents direct bond;

10 Q₂—Q₃ is a single bond;

Q₂ represents CHR² or CO,

wherein

R² represents hydrogen, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy; and

Q₃ represents CHR³,

15 wherein

R³ represents hydrogen, hydroxy, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy,

with the proviso that R² and R³ can not be hydrogen at the same time.

In another embodiment, the urea derivatives of formula (I) are those wherein;

Q₁ represents direct bond;

20 Q₄ represents methylene;

Q₂—Q₃ is a single bond;

Q₂ represents CHR² or CO,

wherein

R^2 represents hydroxy, C_{1-6} alkoxy or C_{1-6} alkanoyloxy;

Q_3 represents CHR^3 ,

wherein

5 R^3 represents hydrogen, hydroxy, C_{1-6} alkoxy or C_{1-6} alkanoyloxy.

In another embodiment, the urea derivatives of formula (I) are those wherein;

Q_1 and Q_4 represent methylene;

Q_2 --- Q_3 is a double bond;

Q_2 represents CR^2 ,

10 wherein

R^2 represents C_{1-6} alkoxy or C_{1-6} alkanoyloxy; and

Q_3 represents CR^3 ,

wherein

R^3 represents hydrogen, C_{1-6} alkoxy or C_{1-6} alkanoyloxy.

15 In another embodiment, the urea derivatives of formula (I) are those wherein;

Q_1 and Q_4 represent methylene;

Q_2 --- Q_3 is a single bond or a double bond;

when Q_2 --- Q_3 is a single bond, Q_2 represents CH_2 and Q_3 represents CHR^3 ,

and when Q_2 --- Q_3 is a double bond, Q_2 represents CH and Q_3 represents CR^3 ,

20 wherein

R^3 represents C_{1-6} alkyl optionally substituted by hydroxy.

..... Preferably, the urea derivatives of formula (I) are those wherein;

n represents an integer of 0 to 1; and

R⁴ represents phenyl optionally substituted with one or more substituents selected from the group consisting of chloro, bromo, fluoro, nitro, methoxy, trifluoromethyl and trifluoromethoxy.

5 More preferably, said urea derivative of the formula (I) is selected from the group consisting of:

N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(6,7-dihydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)urea;

N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(7-hydroxy-6-methoxy-5,6,7,8-tetrahydronaphthalen-1-yl)urea

10 N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(6-hydroxy-7-methoxy-5,6,7,8-tetrahydronaphthalen-1-yl)urea;

4-[(4-Chloro-3-(trifluoromethyl)phenyl)amino]carbonylamino]-2,3-dihydro-1H-inden-2-yl acetate;

15 4-[(4-(Trifluoromethyl)benzyl)amino]carbonylamino]-2,3-dihydro-1H-inden-2-yl acetate;

N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(1-hydroxy-2,3-dihydro-1H-inden-4-yl)urea;

N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(6-hydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)urea; and

N-(6-Hydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)-N'-[4-(trifluoromethyl)benzyl]urea.

20 The urea derivatives of formula (I), their tautomeric and stereoisomeric form, and salts thereof surprisingly show excellent VR1 antagonistic activity. They are, therefore suitable especially for the prophylaxis and treatment of diseases associated with VR1 activity, in particular for the treatment of urinary incontinence, urge urinary incontinence and/or overactive bladder.

25 The compounds of the present invention are also effective for treating or preventing a disease selected from the group consisting of chronic pain, neuropathic pain, postoperative pain, rheumatoid arthritic pain, neuralgia, neuropathies, algesia, nerve injury, ischaemia, neurodegeneration and/or stroke, as well as inflammatory diseases such as asthma and COPD since the diseases also relate to VR1 activity.

The compounds of the present invention are also useful for the treatment and prophylaxis of neuropathic pain, which is a form of pain often associated with herpes zoster and post-herpetic neuralgia, painful diabetic neuropathy, neuropathic low back pain, posttraumatic and postoperative neuralgia, neuralgia due to nerve compression and other neuralgias, phantom pain, complex regional pain syndromes, infectious or parainfectious neuropathies like those associated with HIV infection, pain associated with central nervous system disorders like multiple sclerosis or Parkinson disease or spinal cord injury or traumatic brain injury, and post-stroke pain.

Furthermore, the compounds of the present invention are useful for the treatment of musculoskeletal pain, forms of pain often associated with osteoarthritis or rheumatoid arthritis or other forms of arthritis, and back pain.

In addition, the compounds of the present invention are useful for the treatment of pain associated with cancer, including visceral or neuropathic pain associated with cancer or cancer treatment.

The compounds of the present invention are furthermore useful for the treatment of visceral pain, e.g. pain associated with obstruction of hollow viscus like gallstone colik, pain associated with irritable bowel syndrome, pelvic pain, vulvodynia, orchialgia or prostatodynia, pain associated with inflammatory lesions of joints, skin, muscles or nerves, and orofascial pain and headache, e.g. migraine or tension-type headache.

Further, the present invention provides a medicament, which includes one of the compounds, described above and optionally pharmaceutically acceptable excipients.

Alkyl per se and "alk" and "alkyl" in alkenyl, alkynyl, alkoxy, alkanoyl, alkylamino, alkylamino-carbonyl, alkylaminosulfonyl, alkylsulfonylamino, alkoxy-carbonyl, alkoxy-carbonylamino and alkanoylamino represent a linear or branched alkyl radical having generally 1 to 6, preferably 1 to 4 and particularly preferably 1 to 3 carbon atoms, representing illustratively and preferably methyl, ethyl, n-propyl, isopropyl, tert-butyl, n-pentyl and n-hexyl.

Alkoxy illustratively and preferably represents methoxy, ethoxy, n-propoxy, isopropoxy, tert-butoxy, n-pentoxy and n-hexoxy.

Alkylamino illustratively and preferably represents an alkylamino radical having one or two (independently selected) alkyl substituents, illustratively and preferably representing methylamino, ethylamino, n-propylamino, isopropylamino, tert-butylamino, n-pentylamino, n-hexyl-amino, N,N-dimethylamino, N,N-diethylamino, N-ethyl-N-methylamino, N-methyl-N-n-propylamino, N-

isopropyl-N-n-propylamino, N-t-butyl-N-methylamino, N-ethyl-N-n-pentylamino and N-n-hexyl N-methylamino.

5 Cycloalkyl per se and in cycloalkylamino and in cycloalkylcarbonyl represents a cycloalkyl group having generally 3 to 8 and preferably 5 to 7 carbon atoms, illustratively and preferably representing cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.


10 Heterocyclyl per se and in heterocyclylcarbonyl represents a mono- or polycyclic, preferably mono- or bicyclic, nonaromatic heterocyclic radical having generally 4 to 10 and preferably 5 to 8 ring atoms and up to 3 and preferably up to 2 hetero atoms and/or hetero groups selected from the group consisting of N, O, S, SO and SO₂. The heterocyclyl radicals can be saturated or partially unsaturated. Preference is given to 5- to 8-membered monocyclic saturated heterocyclyl radicals having up to two hetero atoms selected from the group consisting of O, N and S, such as illustratively and preferably tetrahydrofuran-2-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, pyrrolinyl, piperidinyl, morpholinyl, perhydroazepinyl.

EMBODIMENT OF THE INVENTION

15 The compound of the formula (I) of the present invention can be, but not limited to be, prepared by combining various known methods. In some embodiments, one or more of the substituents, such as amino group, carboxyl group, and hydroxyl group of the compounds used as starting materials or intermediates are advantageously protected by a protecting group known to those skilled in the art. Examples of the protecting groups are described in "Protective Groups in Organic Synthesis
20 (3rd Edition)" by Greene and Wuts, John Wiley and Sons, New York 1999.

The compound of the formula (I) of the present invention can be, but not limited to be, prepared by the Method [A], [B], [C], [D], [E] or [F] below.

[Method A]



The compound of the formula (I) (wherein n, Q₁, Q₂, Q₃, Q₄ and R⁴ are the same as defined above) can be prepared by the reaction of the compound of the formula (II) (wherein Q₁, Q₂, Q₃ and Q₄ are the same as defined above) and the compound of the formula (III) (wherein n and R⁴ are the same as defined above).


The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; nitriles such as acetonitrile; amides such as N, N-dimethylformamide (DMF), N, N-dimethylacetamide (DMAC) and N-methylpyrrolidone (NMP); urea such as 1,3-dimethyl-2-imidazolidinone (DMI); sulfoxides such as dimethylsulfoxide (DMSO); and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

The reaction can be carried out in the presence of organic base such as pyridine or triethylamine.

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about room temperature to 100°C. The reaction may be conducted for, usually, 30 minutes to 48 hours and preferably 1 to 24 hours.

The compound (II) and (III) can be prepared by the use of known techniques or are commercially available.

[Method B]




The compound of the formula (I) (wherein n, Q₁, Q₂, Q₃, Q₄ and R⁴ are the same as defined above) can be prepared by reacting the compound of the formula (II) (wherein Q₁, Q₂, Q₃ and Q₄ are the same as defined above) with phosgene, diphosgene, triphosgene, 1,1'-carbonyldiimidazole (CDI), or 1,1'-carbonyldi(1,2,4-triazole)(CDT), and then adding the compound of the formula (IV) (wherein n and R⁴ are the same as defined above) to the reaction mixture.

The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; nitriles such as acetonitrile; amides such as N, N-dimethylformamide (DMF), N, N-dimethylacetamide (DMAC) and N-methylpyrrolidone (NMP); urea such as 1,3-dimethyl-2-imidazolidinone (DMI); sulfoxides such as dimethylsulfoxide (DMSO); and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 20°C to 50°C. The reaction may be conducted for, usually, 30 minutes to 10 hours and preferably 1 to 24 hours.

The compound (IV) is commercially available or can be prepared by the use of known techniques and phosgene, diphosgene, triphosgene, CDI, and CDT are commercially available and .

[Method C]



The compound of the formula (I) (wherein n, Q₁, Q₂, Q₃, Q₄ and R⁴ are the same as defined above) can be prepared by reacting the compound of the formula (II) (wherein Q₁, Q₂, Q₃ and Q₄ are the same as defined above) with the compound of the formula (V) (wherein L₁ represents halogen atom such as chlorine, bromine, or iodine atom) and then adding the compound of the formula (IV) (wherein n and R⁴ are the same as defined above) to the reaction mixture.


The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; nitriles such as acetonitrile; amides such as N, N-dimethylformamide (DMF), N, N-dimethylacetamide (DMAC) and N-methylpyrrolidone (NMP); urea such as 1,3-dimethyl-2-imidazolidinone (DMI); sulfoxides such as dimethylsulfoxide (DMSO); and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 20°C to 50°C. The reaction may be conducted for, usually, 30 minutes to 10 hours and preferably 1 to 24 hours.

The reaction can be advantageously carried out in the presence of a base including, for instance, organic amines such as pyridine, triethylamine and N,N-diisopropylethylamine, dimethylaniline, diethylaniline, 4-dimethylaminopyridine, and others.

The compound (V) is commercially available or can be prepared by the use of known techniques.

[Method D]




The compound of the formula (I) (wherein n, Q₁, Q₂, Q₃, Q₄ and R⁴ are the same as defined above) can be prepared by reacting the compound of the formula (IV) (wherein n and R⁴ are the same as defined above) with phosgene, diphosgene, triphosgene, 1,1'-carbonyldiimidazole (CDI), or 1,1'-

carbonyldi(1,2,4-triazole)(CDT) and then adding the compound of the formula (II) (wherein Q_1 , Q_2 , Q_3 and Q_4 are the same as defined above) to the reaction mixture.

The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; nitriles such as acetonitrile; amides such as N, N-dimethylformamide (DMF), N, N-dimethylacetamide (DMAC) and N-methylpyrrolidone (NMP); urea such as 1,3-dimethyl-2-imidazolidinone (DMI); sulfoxides such as dimethylsulfoxide (DMSO); and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 20°C to 50°C. The reaction may be conducted for, usually, 30 minutes to 10 hours and preferably 1 to 24 hours.

[Method E]



The compound of the formula (I) (wherein n , Q_1 , Q_2 , Q_3 , Q_4 and R^4 are the same as defined above) can be prepared by reacting the compound of the formula (IV) (wherein n and R^4 are the same as defined above) with the compound of the formula (V) (wherein L_1 is the same as defined above) and then adding the compound of the formula (II) (wherein Q_1 , Q_2 , Q_3 and Q_4 are the same as defined above) to the reaction mixture. Q_1 , Q_2 , Q_3 and Q_4 and R^4


The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbons such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; aromatic hydrocarbons such as benzene, toluene and xylene; nitriles such as acetonitrile; amides such as N, N-dimethylformamide (DMF), N, N-dimethylacetamide (DMAC) and N-methylpyrrolidone (NMP); urea such as 1,3-dimethyl-2-imidazolidinone (DMI); sulfoxides such as dimethylsulfoxide

(DMSO); and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 20°C to 50°C. The reaction may be conducted for, usually, 30 minutes to 10 hours and preferably 1 to 24 hours.

The reaction can be advantageously carried out in the presence of a base including, for instance, organic amines such as pyridine, triethylamine and N,N-diisopropylethylamine, dimethylaniline, diethylaniline, 4-dimethylaminopyridine, and others.

[Method F]



The compound of the formula (I-a), (I-b) and (I-c) (wherein n and R^4 are the same as defined above) can be prepared by the following procedures.

In the Step F-1, the compound of the formula (I-a) (wherein n and R^4 are the same as defined above) can be prepared in the similar manner as described in Method [A], [B], [C], [D] or [E] for the preparation of the compound of the formula (I) by using a compound of the formula (VI) instead of the compound of the formula (II).

In the Step F-2, the compound of the formula (I-b) (wherein n and R^4 are the same as defined above) can be prepared by reacting the compound of the formula (I-a) (wherein n and R^4 are the same as defined above) with an acid such as hydrochloric acid.

The reaction may be carried out in a solvent including, for instance, halogenated hydrocarbon such as dichloromethane, chloroform and 1,2-dichloroethane; ethers such as diethyl ether isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; alcohols such as methanol, ethanol; water and others. Optionally, two or more of the solvents selected from the
5 listed above can be mixed and used.

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 20°C to 100°C. The reaction may be conducted for, usually, 30 minutes to 10 hours and preferably 1 to 24 hours.

In the Step F-3, the compound of the formula (I-c) (wherein n and R⁴ are the same as defined
10 above) can be prepared by reacting the compound of the formula (I-b) (wherein n and R⁴ are the same as defined above) with reducing agent such as sodium borohydride or lithium aluminum hydride.

The reaction may be carried out in a solvent including, for instance, ethers such as diethyl ether, isopropyl ether, dioxane and tetrahydrofuran (THF) and 1,2-dimethoxyethane; alcohols such as
15 methanol, ethanol, isopropanol, and others. Optionally, two or more of the solvents selected from the listed above can be mixed and used.

The reaction temperature can be optionally set depending on the compounds to be reacted. The reaction temperature is usually, but not limited to, about 20°C to 50°C. The reaction may be conducted for, usually, 30 minutes to 10 hours and preferably 1 to 24 hours.

20 The compound of the formula (VI) is commercially available or can be prepared by the use of known techniques.

When the compound shown by the formula (I) or a salt thereof has an asymmetric carbon in the structure, their optically active compounds and racemic mixtures are also included in the scope of the present invention.

25 Typical salts of the compound shown by the formula (I) include salts prepared by reaction of the compounds of the present invention with a mineral or organic acid, or an organic or inorganic base. Such salts are known as acid addition and base addition salts, respectively.

Acids to form acid addition salts include inorganic acids such as, without limitation, sulfuric acid, phosphoric acid, hydrochloric acid, hydrobromic acid, hydriodic acid and the like, and organic
30 acids, such as, without limitation, p-toluenesulfonic acid, methanesulfonic acid, oxalic acid, p-

bromophenylsulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like.

Base addition salts include those derived from inorganic bases, such as, without limitation, ammonium hydroxide, alkaline metal hydroxide, alkaline earth metal hydroxides, carbonates, bicarbonates, and the like, and organic bases, such as, without limitation, ethanolamine, triethyl-
5 amine, tris(hydroxymethyl)aminomethane, and the like. Examples of inorganic bases include sodium hydroxide, potassium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like.

The compound of the present invention or a salt thereof, depending on its substituents, may be
10 modified to form lower alkylesters or known other esters; and/or hydrates or other solvates. Those esters, hydrates, and solvates are included in the scope of the present invention.

The compound of the present invention may be administered in oral forms, such as, without limitation normal and enteric coated tablets, capsules, pills, powders, granules, elixirs, tinctures, solution, suspensions, syrups, solid and liquid aerosols and emulsions. They may also be
15 administered in parenteral forms, such as, without limitation, intravenous, intraperitoneal, subcutaneous, intramuscular, and the like forms, well-known to those of ordinary skill in the pharmaceutical arts. The compounds of the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using transdermal delivery systems well-known to those of ordinary skilled in the art.

20 The dosage regimen with the use of the compounds of the present invention is selected by one of ordinary skill in the arts, in view of a variety of factors, including, without limitation, age, weight, sex, and medical condition of the recipient, the severity of the condition to be treated, the route of administration, the level of metabolic and excretory function of the recipient, the dosage form employed, the particular compound and salt thereof employed.

25 The compounds of the present invention are preferably formulated prior to administration together with one or more pharmaceutically-acceptable excipients. Excipients are inert substances such as, without limitation carriers, diluents, flavoring agents, sweeteners, lubricants, solubilizers, suspending agents, binders, tablet disintegrating agents and encapsulating material.

Yet another embodiment of the present invention is pharmaceutical formulation comprising a
30 compound of the invention and one or more pharmaceutically-acceptable excipients that are compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Pharmaceutical formulations of the invention are prepared by combining a therapeutically

effective amount of the compounds of the invention together with one or more pharmaceutically acceptable excipients therefore. In making the compositions of the present invention, the active ingredient may be mixed with a diluent, or enclosed within a carrier, which may be in the form of a capsule, sachet, paper, or other container. The carrier may serve as a diluent, which may be
5 solid, semi-solid, or liquid material which acts as a vehicle, or can be in the form of tablets, pills, powders, lozenges, elixirs, suspensions, emulsions, solutions, syrups, aerosols, ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions and sterile packaged powders.

For oral administration, the active ingredient may be combined with an oral, and non-toxic,
10 pharmaceutically-acceptable carrier, such as, without limitation, lactose, starch, sucrose, glucose, sodium carbonate, mannitol, sorbitol, calcium carbonate, calcium phosphate, calcium sulfate, methyl cellulose, and the like; together with, optionally, disintegrating agents, such as, without limitation, maize, starch, methyl cellulose, agar bentonite, xanthan gum, alginic acid, and the like; and optionally, binding agents, for example, without limitation, gelatin, natural sugars, beta-
15 lactose, corn sweeteners, natural and synthetic gums, acacia, tragacanth, sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like; and, optionally, lubricating agents, for example, without limitation, magnesium stearate, sodium stearate, stearic acid, sodium oleate, sodium benzoate, sodium acetate, sodium chloride, talc, and the like.

In powder forms, the carrier may be a finely divided solid which is in admixture with the finely
20 divided active ingredient. The active ingredient may be mixed with a carrier having binding properties in suitable proportions and compacted in the shape and size desired to produce tablets. The powders and tablets preferably contain from about 1 to about 99 weight percent of the active ingredient which is the novel composition of the present invention. Suitable solid carriers are magnesium carboxymethyl cellulose, low melting waxes, and cocoa butter.

25 Sterile liquid formulations include suspensions, emulsions, syrups and elixirs. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable carriers, such as sterile water, sterile organic solvent, or a mixture of both sterile water and sterile organic solvent.

The active ingredient can also be dissolved in a suitable organic solvent, for example, aqueous propylene glycol. Other compositions can be made by dispersing the finely divided active
30 ingredient in aqueous starch or sodium carboxymethyl cellulose solution or in a suitable oil.

The formulation may be in unit dosage form, which is a physically discrete unit containing a unit dose, suitable for administration in human or other mammals. A unit dosage form can be a

capsule or tablets, or a number of capsules or tablets. A "unit dose" is a predetermined quantity of the active compound of the present invention, calculated to produce the desired therapeutic effect, in association with one or more excipients. The quantity of active ingredient in a unit dose may be varied or adjusted from about 0.1 to about 1000 milligrams or more according to the particular treatment involved.

Typical oral dosages of the present invention, when used for the indicated effects, will range from about 0.01mg /kg/day to about 100 mg/kg/day, preferably from 0.1 mg/kg/day to 30 mg/kg/day, and most preferably from about 0.5 mg/kg/day to about 10 mg/kg/day. In the case of parenteral administration, it has generally proven advantageous to administer quantities of about 0.001 to 100mg /kg/day, preferably from 0.01 mg/kg/day to 1 mg/kg/day. The compounds of the present invention may be administered in a single daily dose, or the total daily dose may be administered in divided doses, two, three, or more times per day. Where delivery is via transdermal forms, of course, administration is continuous.

EXAMPLES

The present invention will be described as a form of examples, but they should by no means be construed as defining the metes and bounds of the present invention.

5 In the examples below, all quantitative data, if not stated otherwise, relate to percentages by weight.

Mass spectra were obtained using electrospray (ES) ionization techniques (micromass Platform LC). Melting points are uncorrected. Liquid Chromatography - Mass spectroscopy (LC-MS) data were recorded on a Micromass Platform LC with Shimadzu Phenomenex ODS column (4.6 mm ϕ \times 30 mm) flushing a mixture of acetonitrile-water (9:1 to 1:9) at 1 ml/min of the flow rate. TLC was performed on a precoated silica gel plate (Merck silica gel 60 F-254). Silica gel (WAKO-gel C-200 (75-150 μ m)) was used for all column chromatography separations. All chemicals were reagent grade and were purchased from Sigma-Aldrich, Wako pure chemical industries, Ltd., Great Britain, Tokyo kasei kogyo Co., Ltd., Nacalai tesque, Inc., Watanabe Chemical Ind. Ltd., Maybridge plc, Lancaster Synthesis Ltd., Merck KGaA, Germany, Kanto Chemical Co., Ltd.

15 ^1H NMR spectra were recorded using either Bruker DRX-300 (300 MHz for ^1H) spectrometer or Bruker 500 UltraShieldTM (500 MHz for ^1H). Chemical shifts are reported in parts per million (ppm) with tetramethylsilane (TMS) as an internal standard at zero ppm. Coupling constant (J) are given in hertz and the abbreviations s, d, t, q, m, and br refer to singlet, doublet, triplet, quartet, multiplet, and broad, respectively. The mass determinations were carried out by MAT95 (Finnigan
20 MAT).

All starting materials are commercially available or can be prepared using methods cited in the literature.

The effect of the present compounds was examined by the following assays and pharmacological tests.

25 [Measurement of capsaicin-induced Ca^{2+} influx in the human VR1-transfected CHO cell line] (Assay 1)

(1) Establishment of the human VR1-CHOluc9aeq cell line

Human vanilloid receptor (hVR1) cDNA was cloned from libraries of axotomized dorsal root ganglia (WO 00/29577). The cloned hVR1 cDNA was constructed with pcDNA3 vector and transfected into a CHOluc9aeq cell line. The cell line contains aequorin and
30

CRE-luciferase reporter genes as read-out signals. The transfectants were cloned by limiting dilution in selection medium (DMEM/F12 medium (Gibco BRL) supplemented with 10% FCS, 1.4 mM Sodium pyruvate, 20 mM HEPES, 0.15% Sodium bicarbonate, 100 U/ml penicillin, 100 µg/ml streptomycin, 2 mM glutamine, non-essential amino acids and 2 mg/ml G418). Ca^{2+} influx was examined in the capsaicin-stimulated clones. A high responder clone was selected and used for further experiments in the project. The human VR1-CHOluc9aeq cells were maintained in the selection medium and passaged every 3-4 days at $1\text{--}2.5 \times 10^5$ cells/flask (75 mm^2).

(2) Measurement of Ca^{2+} influx using FDSS-3000

Human VR1-CHOluc9aeq cells were suspended in a culture medium which is the same as the selection medium except for G418 and seeded at a density of 1,000 cells per well into 384-well plates (black walled clear-base / Nalge Nunc International). Following the culture for 48 hrs the medium was changed to 2 µM Fluo-3 AM (Molecular Probes) and 0.02% Puroic F-127 in assay buffer (Hank's balanced salt solution (HBSS), 17 mM HEPES (pH7.4), 1 mM Probenecid, 0.1% BSA) and the cells were incubated for 60 min at 25°C. After washing twice with assay buffer the cells were incubated with a test compound or vehicle for 20 min at 25°C. Mobilization of cytoplasmic Ca^{2+} was measured by FDSS-3000 ($\lambda_{\text{ex}}=488\text{nm}$, $\lambda_{\text{em}}=540\text{nm}$ / Hamamatsu Photonics) for 60 sec after the stimulation with 10 nM capsaicin. Integral R was calculated and compared with controls.

20 [Measurement of the capsaicin-induced Ca^{2+} influx in primary cultured rat dorsal root ganglia neurons] (Assay 2)

(1) Preparation of rat dorsal root ganglia neurons

New born Wister rats (5-11 days) were sacrificed and dorsal root ganglia (DRG) was removed. DRG was incubated with 0.1% trypsin (Gibco BRL) in PBS(-) (Gibco BRL) for 30 min at 37°C, then a half volume of fetal calf serum (FCS) was added and the cells were spun down. The DRG neuron cells were resuspended in Ham F12/5% FCS/5% horse serum (Gibco BRL) and dispersed by repeated pipetting and passing through 70 µm mesh (Falcon). The culture plate was incubated for 3 hours at 37°C to remove contaminating Schwann cells. Non-adherent cells were recovered and further cultured in laminin-coated 384 well plates (Nunc) at 1×10^4 cells/50 µl/well for 2 days in the presence of 50 ng/ml recombinant rat NGF-(Sigma)-and 50-µM 5-fluorodeoxyuridine (Sigma).

(2) Ca^{2+} mobilization assay

DRG neuron cells were washed twice with HBSS supplemented with 17 mM HEPES (pH 7.4) and 0.1% BSA. After incubating with 2 μM fluo-3AM (Molecular Probe), 0.02% PF127 (Gibco BRL) and 1 mM probenecid (Sigma) for 40 min at 37°C, cells were washed 3 times. The cells were incubated with VR1 antagonists or vehicle (dimethylsulfoxide) and then with 1 μM capsaicin in FDSS-6000 (λ_{ex} =480nm, λ_{em} =520nm / Hamamatsu Photonics). The fluorescence changes at 480nm were monitored for 2.5 min. Integral R was calculated and compared with controls.

[Organ bath assay to measure the capsaicin-induced bladder contraction] (Assay 3)

Male Wistar rats (10 week old) were anesthetized with ether and sacrificed by dislocating the necks. The whole urinary bladder was excised and placed in oxygenated Modified Krebs-Henseleit solution (pH 7.4) of the following composition (112mM NaCl, 5.9 mM KCl, 1.2 mM MgCl_2 , 1.2 mM NaH_2PO_4 , 2 mM CaCl_2 , 2.5 mM NaHCO_3 , 12 mM glucose). Contractile responses of the urinary bladder were studied as described previously [Maggi CA et al: Br.J.Pharmacol. 108: 801-805, 1993]. Isometric tension was recorded under a load of 1 g using longitudinal strips of rat detrusor muscle. Bladder strips were equilibrated for 60 min before each stimulation. Contractile response to 80 mM KCl was determined at 15 min intervals until reproducible responses were obtained. The response to KCl was used as an internal standard to evaluate the maximal response to capsaicin. The effects of the compounds were investigated by incubating the strips with compounds for 30 min prior to the stimulation with 1 μM capsaicin (vehicle: 80% saline, 10% EtOH, and 10% Tween 80). One of the preparations made from the same animal was served as a control while the others were used for evaluating compounds. Ratio of each capsaicin-induced contraction to the internal standard (i.e. KCl-induced contraction) was calculated and the effects of the test compounds on the capsaicin-induced contraction were evaluated.

[Measurement of Ca^{2+} influx in the human P2X1-transfected CHO cell line](1) Preparation of the human P2X1-transfected CHO_{luc9aeq} cell line

Human P2X1-transfected CHO_{luc9aeq} cell line was established and maintained in Dulbecco's modified Eagle's medium (DMEM/F12) supplemented with 7.5% FCS, 20 mM HEPES-KOH (pH 7.4), 1.4 mM sodium pyruvate, 100 U/ml penicillin, 100 $\mu\text{g}/\text{ml}$ streptomycin, 2 mM glutamine (Gibco BRL) and 0.5 Units/ml apyrase (grade I, Sigma). The suspended cells were seeded in each well of 384-well optical bottom black plates

(Nalge Nunc International) at 3×10^3 / 50 μ l / well. The cells were cultured for following 48 hrs to adhere to the plates.

(2) Measurement of the intracellular Ca^{2+} levels

P2X1 receptor agonist-mediated increases in cytosolic Ca^{2+} levels were measured using a fluorescent Ca^{2+} chelating dye, Fluo-3 AM (Molecular Probes). The plate-attached cells were washed twice with washing buffer (HBSS, 17 mM HEPES-KOH (pH 7.4), 0.1% BSA and 0.5 units/ml apyrase), and incubated in 40 μ l of loading buffer (1 μ M Fluo-3 AM, 1 mM probenecid, 1 μ M cyclosporin A, 0.01% pluronic (Molecular Probes) in washing buffer) for 1 hour in a dark place. The plates were washed twice with 40 μ l washing buffer and 35 μ l of washing buffer were added in each well with 5 μ l of test compounds or 2',3'-*o*-(2,4,6-trinitrophenyl) adenosine 5'-triphosphate (Molecular Probes) as a reference. After further incubation for 10 minutes in dark 200 nM α , β -methylene ATP agonist was added to initiate the Ca^{2+} mobilization. Fluorescence intensity was measured by FDSS-6000 (λ_{ex} =410nm, λ_{em} =510nm / Hamamatsu Photonics) at 250 msec intervals. Integral ratios were calculated from the data and compared with that of a control.

[Measurement of capsaicin-induced bladder contraction in anesthetized rats] (Assay 4)

(1) Animals

Female Sprague-Dawley rats (200~250 g / Charles River Japan) were used.

(2) Catheter implantation

Rats were anesthetized by intraperitoneal administration of urethane (Sigma) at 1.2 g/kg. The abdomen was opened through a midline incision, and a polyethylene catheter (BECTON DICKINSON, PE50) was implanted into the bladder through the dome. In parallel, the inguinal region was incised, and a polyethylene catheter (Hibiki, size 5) filled with 2 IU / ml of heparin (Novo Heparin, Aventis Pharma) in saline (Otsuka) was inserted into a common iliac artery.

(3) Cystometric investigation

The bladder catheter was connected via T-tube to a pressure transducer (Viggo-Spectramed Pte Ltd, DT-XXAD) and a microinjection pump (TERUMO). Saline was infused at room temperature into the bladder at a rate of 2.4 ml/hr. Intravesical pressure was recorded continuously on a chart pen recorder (Yokogawa). At least three

reproducible micturition cycles, corresponding to a 20-minute period, were recorded before a test compound administration and used as baseline values.

(4) Administration of test compounds and stimulation of bladder with capsaicin

5 The saline infusion was stopped before administering compounds. A testing compound dissolved in the mixture of ethanol, Tween 80 (ICN Biomedicals Inc.) and saline (1 : 1 : 8, v/v/v) was administered intraarterially at 10 mg/kg. 2min after the administration of the compound 10 µg of capsaicin (Nacalai Tesque) dissolved in ethanol was administered intraarterially.

(5) Analysis of cystometry parameters

10 Relative increases in the capsaicin-induced intravesical pressure were analyzed from the cystometry data. The capsaicin-induced bladder pressures were compared with the maximum bladder pressure during micturition without the capsaicin stimulation. The testing compounds-mediated inhibition of the increased bladder pressures was evaluated using Student's t-test. A probability level less than 5% was accepted as significant
15 difference.

[Measurement of over active bladder in anesthetized cystitis rats] (Assay 5)

(1) Animals

Female Sprague-Dawley rats (180~250 g / Charles River Japan) were used. Cyclophosphamide (CYP) dissolved in saline was administered intraperitoneally at 150 mg/kg
20 48 hours before experiment.

(2) Catheter implantation

25 Rats were anesthetized by intraperitoneal administration of urethane (Sigma) at 1.25 g/kg. The abdomen was opened through a midline incision, and a polyethylene catheter (BECTON DICKINSON, PE50) was implanted into the bladder through the dome. In parallel, the inguinal region was incised, and a polyethylene catheter (BECTON DICKINSON, PE50) filled with saline (Otsuka) was inserted into a femoral vein. After the bladder was emptied, the rats were left for 1 hour for recovery from the operation.

(3) Cystometric investigation

The bladder catheter was connected via T-tube to a pressure transducer (Viggo-Spectramed Pte Ltd, DT-XXAD) and a microinjection pump (TERUMO). Saline was infused at room temperature into the bladder at a rate of 3.6 ml/hr for 20 min. Intravesical pressure was recorded continuously on a chart pen recorder (Yokogawa). At least three reproducible micturition cycles, corresponding to a 20-minute period, were recorded before a test compound administration.

(4) Administration of test compounds

A testing compound dissolved in the mixture of ethanol, Tween 80 (ICN Biomedicals Inc.) and saline (1 : 1 : 8, v/v/v) was administered intravenously at 0.05 mg/kg, 0.5 mg/kg or 5 mg/kg. 3min after the administration of the compound, saline (Nacalai Tesque) was infused at room temperature into the bladder at a rate of 3.6 ml/hr.

(5) Analysis of cystometry parameters

The cystometry parameters were analyzed as described previously [Lecci A et al: Eur. J. Pharmacol. 259: 129-135, 1994]. The micturition frequency calculated from micturition interval and the bladder capacity calculated from a volume of infused saline until the first micturition were analyzed from the cystometry data. The testing compounds-mediated inhibition of the frequency and the testing compounds-mediated increase of bladder capacity were evaluated using unpaired Student's t-test. A probability levels less than 5% was accepted as significant difference. Data were analyzed as the mean \pm SEM from 4 – 7 rats.

[Measurement of Acute Pain]

Acute pain is measured on a hot plate mainly in rats. Two variants of hot plate testing are used: In the classical variant animals are put on a hot surface (52 to 56°C) and the latency time is measured until the animals show nociceptive behavior, such as stepping or foot licking. The other variant is an increasing temperature hot plate where the experimental animals are put on a surface of neutral temperature. Subsequently this surface is slowly but constantly heated until the animals begin to lick a hind paw. The temperature which is reached when hind paw licking begins is a measure for pain threshold.

Compounds are tested against a vehicle treated control group. Substance application is performed at different time points via different application routes (i.v., i.p., p.o., i.t., i.c.v., s.c., intradermal, transdermal) prior to pain testing.

[Measurement of Persistent Pain]

- 5 Persistent pain is measured with the formalin or capsaicin test, mainly in rats. A solution of 1 to 5% formalin or 10 to 100 µg capsaicin is injected into one hind paw of the experimental animal. After formalin or capsaicin application the animals show nociceptive reactions like flinching, licking and biting of the affected paw. The number of nociceptive reactions within a time frame of up to 90 minutes is a measure for intensity of pain.
- 10 Compounds are tested against a vehicle treated control group. Substance application is performed at different time points via different application routes (i.v., i.p., p.o., i.t., i.c.v., s.c., intradermal, transdermal) prior to formalin or capsaicin administration.

[Measurement of Neuropathic Pain]

- Neuropathic pain is induced by different variants of unilateral sciatic nerve injury mainly in rats.
- 15 The operation is performed under anesthesia. The first variant of sciatic nerve injury is produced by placing loosely constrictive ligatures around the common sciatic nerve (Bennett and Xie, Pain 33 (1988): 87-107). The second variant is the tight ligation of about the half of the diameter of the common sciatic nerve (Seltzer et al., Pain 43 (1990): 205-218). In the next variant, a group of models is used in which tight ligations or transections are made of either the L5 and L6 spinal
- 20 nerves, or the L5 spinal nerve only (KIM SH; CHUNG JM, AN EXPERIMENTAL-MODEL FOR PERIPHERAL NEUROPATHY PRODUCED BY SEGMENTAL SPINAL NERVE LIGATION IN THE RA, PAIN 50 (3) (1992): 355-363). The fourth variant involves an axotomy of two of the three terminal branches of the sciatic nerve (tibial and common peroneal nerves) leaving the remaining sural nerve intact whereas the last variant comprises the axotomy of only the tibial
- 25 branch leaving the sural and common nerves uninjured. Control animals are treated with a sham operation.

- Postoperatively, the nerve injured animals develop a chronic mechanical allodynia, cold allodynia, as well as a thermal hyperalgesia. Mechanical allodynia is measured by means of a pressure transducer (electronic von Frey Anesthesiometer, IITC Inc.-Life Science Instruments, Woodland Hills, SA, USA; Electronic von Frey System, Somedic Sales AB, Hörby, Sweden). Thermal hyperalgesia is measured by means of a radiant heat source (Plantar Test, Ugo Basile, Comerio, Italy), or by means of a cold plate of 5 to 10 °C where the nocifensive reactions of the affected
- 30

hind paw are counted as a measure of pain intensity. A further test for cold induced pain is the counting of nocifensive reactions, or duration of nocifensive responses after plantar administration of acetone to the affected hind limb. Chronic pain in general is assessed by registering the circadian rhythms in activity (Surjo and Arndt, Universität zu Köln, Cologne, Germany), and by scoring differences in gait (foot print patterns; FOOTPRINTS program, Klapdor et al., 1997. A low cost method to analyse footprint patterns. J. Neurosci. Methods 75, 49-54).

Compounds are tested against sham operated and vehicle treated control groups. Substance application is performed at different time points via different application routes (i.v., i.p., p.o., i.t., i.c.v., s.c., intradermal, transdermal) prior to pain testing.

10 [Measurement of Inflammatory Pain]

Inflammatory pain is induced mainly in rats by injection of 0.75 mg carrageenan or complete Freund's adjuvant into one hind paw. The animals develop an edema with mechanical allodynia as well as thermal hyperalgesia. Mechanical allodynia is measured by means of a pressure transducer (electronic von Frey Anesthesiometer, IITC Inc.-Life Science Instruments, Woodland Hills, SA, USA). Thermal hyperalgesia is measured by means of a radiant heat source (Plantar Test, Ugo Basile, Comerio, Italy, Paw thermal stimulator, G. Ozaki, University of California, USA). For edema measurement two methods are being used. In the first method, the animals are sacrificed and the affected hindpaws sectioned and weighed. The second method comprises differences in paw volume by measuring water displacement in a plethysmometer (Ugo Basile, Comerio, Italy).

20 Compounds are tested against uninflamed as well as vehicle treated control groups. Substance application is performed at different time points via different application routes (i.v., i.p., p.o., i.t., i.c.v., s.c., intradermal, transdermal) prior to pain testing.

[Measurement of Diabetic Neuropathic Pain]

25 Rats treated with a single intraperitoneal injection of 50 to 80 mg/kg streptozotocin develop a profound hyperglycemia and mechanical allodynia within 1 to 3 weeks. Mechanical allodynia is measured by means of a pressure transducer (electronic von Frey Anesthesiometer, IITC Inc.-Life Science Instruments, Woodland Hills, SA, USA).

Compounds are tested against diabetic and non-diabetic vehicle treated control groups. Substance application is performed at different time points via different application routes (i.v., i.p., p.o., i.t., i.c.v., s.c., intradermal, transdermal) prior to pain testing.

30


Results in capsaicin-induced Ca^{2+} influx in the human VR1-transfected CHO cell line (Assay 1 are shown in Examples and tables of the Examples below. For practical reasons, the compounds are grouped in four classes of activity as follows:

$$\text{IC}_{50} = \text{A} (< \text{or} =) 0.1 \mu\text{M} < \text{B} (< \text{or} =) 0.5 \mu\text{M} < \text{C} (< \text{or} =) 1 \mu\text{M} < \text{D}$$

- 5 The compounds of the present invention also show excellent selectivity, and strong activity in other assays 2-5 and assays for pain described above.

Preparing method of starting compounds

4-Amino-2,3-dihydro-1H-inden-2-yl acetate



To a solution of 2-nitrobenzyl bromide (1.00 g, 4.63 mmol) and diethyl malonate (0.741 g, 4.63 mmol) in 30 ml of hexane was added potassium carbonate (0.640 g, 4.63 mmol) and 18-Crown-6 (0.012 g, 0.05 mmol). After stirred at 80 °C for 18 hours, the mixture was diluted with

water and was extracted with ethyl acetate. The organic layer was washed with water, then with brine, and concentrated under reduced pressure to obtain crude diethyl (2-nitrobenzyl)malonate.

A solution of crude diethyl (2-nitrobenzyl)malonate in 6N aqueous HCl (15 ml) and acetic acid (15 ml) was stirred at refluxing temperature for 48 hours. After cooled to ambient temperature, the mixture was concentrated under reduced pressure. To the residue was added 10% aqueous NaOH solution and washed with ethyl acetate. The aqueous layer was acidified with aqueous HCl solution, and the mixture was extracted with ethyl acetate. The organic layer was dried over MgSO₄, filtered, and concentrated under reduced pressure to obtain 3-(2-nitrophenyl)propanoic acid.

¹H NMR (CDCl₃) δ 2.79 (t, *J* = 7.6 Hz, 2H), 3.24 (t, *J* = 7.6 Hz, 2H), 7.38-7.44 (m, 2H), 7.55 (dt, *J* = 7.6, 1.6 Hz, 1H), 7.96 (dd, *J* = 7.6, 1.6 Hz, 1H).

A solution of 3-(2-nitrophenyl)propanoic acid (1.20 g, 6.15 mmol) and thionyl chloride (0.878 g, 7.38 mmol) in dichloromethane (5 ml) was stirred and heated to reflux for 2 hours. The mixture was concentrated under reduced pressure to obtain 3-(2-nitrophenyl)propanoyl chloride. To the obtained crude 3-(2-nitrophenyl)propanoyl chloride (1.31 g, 6.15 mmol) was added CS₂, and aluminum trichloride (1.07 g, 8.0 mmol) was added portionwise at 0°C. The mixture was stirred at 70°C for 3 hours, and after cooled to ambient temperature, water was added and extracted with ethyl acetate. The organic layer was dried over MgSO₄, filtered, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane:ethylacetate 10:1) to afford 4-nitroindan-1-one (0.44 g).

¹H NMR (CDCl₃) δ 2.79-2.82 (m, 2H), 3.64-3.66 (m, 2H), 7.62 (t, *J* = 7.9 Hz, 1H), 8.09 (d, *J* = 7.6 Hz, 1H), 8.47 (d, *J* = 8.2 Hz, 1H).

To a solution of 4-nitroindan-1-one (0.381 g, 2.15 mmol) in ethanol (5 ml) was added sodium borohydride (0.048 g, 1.29 mmol) at 0 °C, and the mixture was stirred at room temperature for 3 hours. Aqueous solution of ammonium chloride was added to the mixture, and extracted with ethyl acetate. The organic layer was dried over MgSO₄, filtered, and concentrated under reduced pressure to obtain 4-nitroindan-1-ol. ¹H NMR (CDCl₃) δ 1.90 (d, *J* = 6.5 Hz, 1H), 2.00-2.07 (m, 1H), 2.56-2.63 (m, 1H), 3.25-3.33 (m, 1H), 3.54-3.60 (m, 1H), 5.30-5.35 (m, 1H), 7.44 (t, *J* = 8.2 Hz, 1H), 7.72 (d, *J* = 7.6 Hz, 1H), 8.12 (d, *J* = 8.2 Hz, 1H).

A solution of 4-nitroindan-1-ol (0.385 g, 2.15 mmol) and *p*-toluenesulfonic acid (5.0 mg, 0.03 mmol) in toluene (30 ml) was stirred and heated to reflux for 16 hours. After cooled to ambient temperature, the mixture was washed with aqueous sodium bicarbonate solution. The

organic layer was dried over MgSO_4 , filtered, and concentrated under reduced pressure. The obtained residue was purified by preparatory TLC (hexane:ethylacetate 3:1) to afford 7-nitro-1*H*-indene (0.289 g).

^1H NMR (CDCl_3) δ 3.94 (s, 2H), 6.75 (dt, $J = 5.7, 1.9$ Hz, 1H), 6.93 (dt, $J = 5.7, 1.6$ Hz, 1H), 7.4 (t, $J = 8.2$ Hz, 1H), 7.68 (d, $J = 7.6$ Hz, 1H), 8.05 (d, $J = 8.2$ Hz, 1H).

To a solution of 2,3-dimethyl-2-butene (21.5 mg, 0.31 mmol) in THF (2 ml) at 0°C was added borane-THF (0.307 ml, 0.31 mmol) dropwise. After stirred for 1 hour at 0°C , 7-nitro-1*H*-indene (45.0 mg, 0.28 mmol) in THF (5 ml) was added dropwise, and the mixture was stirred for 2 hours at ambient temperature. The mixture was cooled to 0°C , and water (0.15 ml), 4*N* aqueous sodium hydroxide (0.45 ml), and 30% H_2O_2 (0.45 ml) were added. The mixture was then warmed to room temperature and poured into water, extracted with ethyl acetate and washed with brine. The organic layer was dried over MgSO_4 , filtered, and concentrated under reduced pressure. To the obtained mixture in toluene (1 ml) was added acetic anhydride (40.8 mg, 0.40 mmol) and pyridine (0.4 ml), and then stirred for 16 hours at room temperature. The mixture was concentrated under reduced pressure, and the obtained residue was purified by preparatory TLC (hexane:ethylacetate 2:1) to obtain 4-nitro-2,3-dihydro-1*H*-inden-2-yl acetate (16.0 mg).


^1H NMR (CDCl_3) δ 2.03 (s, 3H), 3.12 (dd, $J = 17.5, 1.6$ Hz, 1H), 3.40 (dd, $J = 17.5, 6.3$ Hz, 1H), 3.60 (dd, $J = 19.2, 2.2$ Hz, 1H), 3.74 (dd, $J = 19.2, 6.6$ Hz, 1H), 5.58-5.62 (m, 1H), 7.39 (t, $J = 7.9$ Hz, 1H), 7.54 (d, $J = 7.3$ Hz, 1H), 8.06 (d, $J = 8.2$ Hz, 1H).

To a mixture of 4-nitro-2,3-dihydro-1*H*-inden-2-yl acetate (100 mg, 0.45 mmol) and ammonium chloride (100 mg) in ethanol (6 ml) and water (3 ml) was added iron powder (300 mg) portionwise at room temperature. The mixture was stirred at 90°C for 1 hour, and after cooled to room temperature, the mixture was diluted with ethylacetate. The mixture was filtered through a pad of celite, and the filtrate was washed with brine, dried over MgSO_4 , filtered, and concentrated under reduced pressure to obtain 4-amino-2,3-dihydro-1*H*-inden-2-yl acetate.

^1H NMR (CDCl_3) δ 2.03 (s, 3H), 2.81 (dd, $J = 16.4, 2.8$ Hz, 1H), 3.00 (dd, $J = 16.7, 2.8$ Hz, 1H), 3.14 (dd, $J = 16.4, 6.6$ Hz, 1H), 3.29 (dd, $J = 16.7, 6.6$ Hz, 1H), 3.58 (br.s, 2H), 5.51-5.56 (m, 1H), 6.54 (d, $J = 7.9$ Hz, 1H), 6.69 (d, $J = 7.3$ Hz, 1H), 7.04 (t, $J = 7.9$ Hz, 1H).

Example 1-1

4-[[[4-Chloro-3-(trifluoromethyl)phenyl]amino}carbonyl]amino]-2,3-dihydro-1H-inden-2-yl acetate



- 5 A mixture of 4-amino-2,3-dihydro-1H-inden-2-yl acetate (86.4 mg, 0.45 mmol) and 4-chloro-3-trifluoromethylphenyl isocyanate (110 mg, 0.50 mmol) in 1,4-dioxane (2 m) was stirred at 50°C for 15 hours. The mixture was concentrated under reduced pressure, and to the obtained residue was added diisopropyl ether. The precipitate was collected to afford 4-[[[4-chloro-3-(trifluoromethyl)phenyl]amino}carbonyl]amino]-2,3-dihydro-1H-inden-2-yl acetate (128 mg).
- 10 ¹H NMR (DMSO-*d*₆) δ 1.98 (s, 3H), 2.91 (ddd, *J*= 19.6, 17.1, 1.9 Hz, 2H), 3.21-3.30 (m, 2H), 5.40-5.45 (m, 1H), 6.96 (d, *J*= 7.3 Hz, 1H), 7.15 (t, *J*= 7.9 Hz, 1H), 7.62 (s, 2H), 7.71 (d, *J*= 8.2 Hz, 1H), 8.10 (s, 1H), 8.25 (s, 1H), 9.34 (s, 1H);

Molecular weight : 412.80



MS (M+H): 413

- 15 Mp 207-209°C;

Activity class: C


In the similar manner as described in Example 1-1, compounds in Example 1-2 to 1-3 as shown in Table 1 were synthesized.

Table 1

example #	structure	M.W.	MS (M+1)	MP	activity class
1-2		392,38	393	166-168	A
1-3		370,76	371	221-223	A

Starting material

(6-Ethoxy-5,8-dihydronaphthalen-1-yl)amine



A mixture of 5-amino-2-naphthol (4.78 g, 30.0 mmol), benzaldehyde (3.50 g, 33.0 mmol), and magnesium sulfate (10.0 g) in THF (100 ml) was heated to reflux for 16 hours. After cooled to

ambient temperature, the mixture was filtered through a pad of celite and the filtrate was concentrated under reduced pressure. The obtained residue was recrystallized with diethylether to afford 5-[[phenylmethylene]amino]-2-naphthol (7.40 g).

¹H NMR (CDCl₃) δ 5.06 (br.s, 1H), 6.92 (d, *J* = 6.6 Hz, 1H), 7.10-7.17 (m, 2H), 7.42-7.49 (dd, *J* = 6.6 Hz, 1H), 7.45-7.55 (m, 4H), 8.00-8.02 (m, 2H), 8.27 (d, *J* = 9.0 Hz, 1H), 8.56 (s, 1H)

Molecular weight : 247.30

MS (M+H): 248

To a solution of 5-[(phenylmethylene)amino]-2-naphthol (2.00 g, 8.09 mmol) in DMF (50 ml) was added ethyl iodide (1.39 g, 8.90 mmol) at room temperature and stirred at 50°C for 2 hours. After cooled to ambient temperature, water was added and the mixture was extracted with ethyl acetate. The organic layer was washed with water and brine, dried over MgSO₄, filtered, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane:ethylacetate 15:1) to afford (6-ethoxy-1-naphthyl)(phenylmethylene)-amine (1.54 g).

¹H NMR (CDCl₃) : δ 1.49 (3H, t, *J* = 6.8 Hz), 4.17 (2H, q, *J* = 6.8 Hz), 6.91 (1H, dd, *J* = 1.1, 7.5 Hz), 7.14-7.18 (2H, m), 7.41 (1H, dd, *J* = 7.2, 7.2 Hz), 7.50-7.61 (4H, m), 7.99-8.03 (2H, m), 8.25 (1H, d, *J* = 8.7 Hz), 8.55 (1H, s);

Molecular weight : 275.35

MS (M+H): 276

A mixture of (6-ethoxy-1-naphthyl)(phenylmethylene)amine (0.600 g, 2.18 mmol) and Pd/C (0.900 g) in ethyl acetate (15 ml) was stirred under argon at room temperature for 48 hours. The mixture was filtered through a pad of celite, and the filtrate was concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane:ethylacetate 4:1) to provide (6-ethoxy-1-naphthyl)amine (2.78 g).

¹H NMR (CDCl₃) : δ 1.46 (3H, t, *J* = 6.8 Hz), 4.06 (2H, brs), 4.13 (2H, q, *J* = 6.8 Hz), 6.62 (1H, dd, *J* = 1.5, 6.8 Hz), 7.08-7.12 (2H, m), 7.16-7.25 (2H, m), 7.70 (1H, d, *J* = 9.8 Hz)

Molecular weight : 187.24

MS (M+H): 188

To a mixture of (6-ethoxy-1-naphthyl)amine (300 mg, 1.60 mmol) and tert-buthanol (641 mg, 8.65 mmol) in THF (4 ml) and liquid ammonia (55 ml) at -78°C was added lithium (96.8 mg, 13.94 mmol) portionwise. After the mixture was stirred for 30 minutes at -78°C , methanol (9 ml) and water were added. Ammonia was removed at room temperature, and the resulted mixture was
5 extracted with ethyl acetate. The organic layer was dried over MgSO_4 , filtered, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane:ethylacetate 4:1) to afford (6-ethoxy-5,8-dihydronaphthalen-1-yl)amine (248 mg).


^1H NMR (CDCl_3) : δ 1.33 (3H, t, $J = 6.8$ Hz), 3.17 (1H, dd, $J = 3.4, 5.1$ Hz), 3.20 (1H, dd, $J = 3.4, 5.1$ Hz), 3.42 (1H, d, $J = 5.1$ Hz), 3.43 (1H, d, $J = 5.1$ Hz), 3.57 (2H, brs), 3.81 (2H, q, $J = 6.8$ Hz),
10 4.77 (1H, t, $J = 3.4$ Hz), 6.52 (1H, d, $J = 7.9$ Hz), 6.58 (1H, d, $J = 7.5$ Hz), 6.98 (1H, dd, $J = 7.5, 7.9$ Hz).

Molecular weight : 189.26

MS ($\text{M}+\text{H}$): 190

Example 2-1

15 N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(6-ethoxy-5,8-dihydronaphthalen-1-yl)urea



Next, to a solution of (6-ethoxy-5,8-dihydronaphthalen-1-yl)amine (108 mg, 0.57 mmol) in THF (5. ml) was added 4-chloro-3-trifluoromethyl isocyanate (139 mg, 0.63 mmol), and the mixture was stirred for 13 hours. Saturated aqueous solution of sodium carbonate was added and the mixture was extracted with ethylacetate. The organic layer was dried over MgSO_4 , filtered, and concentrated under reduced pressure to obtain N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(6-ethoxy-5,8-dihydronaphthalen-1-yl)urea (234 mg).
20

¹H NMR (DMSO-*d*₆): δ 1.26 (3H, t, *J* = 6.8 Hz), 3.29-3.38 (4H, m), 3.80 (2H, q, *J* = 6.8 Hz), 6.91 (1H, d, *J* = 7.5 Hz), 7.14 (1H, dd, *J* = 7.5, 7.9 Hz), 7.59-7.61 (2H, m), 8.01 (1H, s), 8.10 (1H, s), 9.45 (1H, s)

Molecular weight : 410.82


5 MS (M+H): 411

Mp 216°C;

Activity class: B

Example 2-2

N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(6-oxo-5,6,7,8-tetrahydronaphthalen-1-yl)urea



To a solution of N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(6-ethoxy-5,8-dihydronaphthalen-1-yl)urea (50.0 mg, 0.12 mmol) was added aqueous 1N HCl solution at room temperature. After stirred for 20 minutes, saturated aqueous solution of sodium carbonate was added and the mixture was extracted with ethylacetate. The organic layer was dried over MgSO₄, filtered, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (hexane:ethylacetate 1:2) to afford N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(6-oxo-5,6,7,8-tetrahydronaphthalen-1-yl)urea (41.7 mg).

15

¹H NMR (Acetone-*d*₆) : δ 2.44 (2H, t, *J* = 6.4 Hz), 3.06 (2H, t, *J* = 6.4 Hz), 3.57 (2H, s), 6.98 (1H, d, *J* = 7.2 Hz), 7.19 (1H, dd, *J* = 7.2, 7.5 Hz), 7.51-7.52 (2H, m), 7.74 (1H, dd, *J* = 2.6, 8.7 Hz), 7.87 (1H, brs), 7.14 (1H, d, *J* = 2.6 Hz), 8.69 (1H, brs);

20

Molecular weight : 382.77


MS (M+H): 383

Mp 219°C;

Activity class: A

Example 2-3

N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(6-hydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)urea



To a solution of N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(6-oxo-5,6,7,8-tetrahydronaphthalen-1-yl)urea (70.0 mg, 0.18 mmol) in methanol (3 ml) was added sodium borohydride (7.61 mg, 0.20 mmol) at 0 °C. After stirred for 30 minutes, the mixture was concentrated under reduced pressure and water was added. The mixture was extracted with ethylacetate, and the organic layer
10 was dried over MgSO₄, filtered, and concentrated under reduced pressure to obtain N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(6-hydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)urea (70.0 mg).

¹H NMR (Acetone-*d*₆) : δ 1.75 (1H, m), 2.04 (1H, m), 2.59-3.04 (4H, m), 4.02 (1H, m), 6.84 (1H, d, *J* = 7.2 Hz), 7.09 (1H, dd, *J* = 7.2, 7.5 Hz), 7.50-7.53 (2H, m), 7.67 (1H, d, *J* = 7.5 Hz), 7.72 (1H, dd, *J* = 2.6, 8.7 Hz), 8.13 (1H, d, *J* = 2.6 Hz), 8.77 (1H, s);

15 Molecular weight : 384.79






MS (M+H): 385

Mp 216°C

Activity class: A

In the similar manner as described in Example 2-1, 2-2, or 2-3, compounds in Example 2-4 to 2-9
20 as shown in Table 2 were synthesized.


Table 2

example #	structure	M.W.	MS (M+1)	MP	activity class
2-4		390,41	391	235	C
2-5		362,35	363	221	A
2-6		364,37	365	205	A
2-7		400,79	400	201	A
2-8		414,82	414	220-222	B

example #	structure	M.W.	MS (M+1)	MP	activity class
2-9		414,82	414	112-127	A

Starting material

(7-Methyl-5,6,7,8-tetrahydronaphthalen-1-yl)amine



- 5 A mixture of 8-amino-3,4-dihydronaphthalen-2(1H)-one (1.61 g, 9.99 mmol), benzyl bromide (1.88 g, 11.0 mmol), and potassium carbonate (2.07 g, 15.0 mmol) in acetone (50 mL) was stirred at refluxing temperature for 16 hours. After the mixture was cooled to ambient temperature, it was filtered through a pad of Celite, and the filtrate was concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (eluent: ethylacetate / hexane
10 = 1 / 10) to provide 8-(benzylamino)-3,4-dihydronaphthalen-2(1H)-one (1.87 g).

¹H NMR (CDCl₃): δ 2.65 (dd, *J* = 12.9 Hz, 6.6 Hz, 2H), 3.10 (dd, *J* = 12.9 Hz, 6.6 Hz, 2H), 3.28 (s, 2H), 3.71 (brs, 1H), 4.31 (s, 2H), 6.55 (d, *J* = 8.1 Hz, 1H), 6.60 (d, *J* = 8.1 Hz, 1H), 7.12 (t, *J* = 8.1 Hz, 1H), 7.23 – 7.40 (m, 5H);

Molecular weight: 251.33

- 15 MS (M+H): 252

To a suspension of methyltriphenylphosphonium iodide (2.12 g, 5.25 mmol) in tetrahydrofuran (100 ml) was added sodium *tert*-butoxide (0.56 g, 5.83 mmol) at 0°C. After the mixture was stirred for 30 minutes, a solution of 8-(benzylamino)-3,4-dihydronaphthalen-2(1H)-one (0.66 g, 2.63 mmol) in tetrahydrofuran (10 ml) was added at room temperature and then stirred at 100 °C for 13 hours. The mixture was cooled to ambient temperature and was poured into water. The mixture was extracted with ethylacetate, and the organic layer was dried over Na₂SO₄, filtered, and concentrated under reduced pressure to obtain N-benzyl-7-methylene-5,6,7,8-tetrahydronaphthalen-1-amine (0.367 g).

¹H NMR (CDCl₃): δ 2.45 (dd, *J* = 12.9 Hz, 6.6 Hz, 2H), 2.86 (dd, *J* = 12.9 Hz, 6.6 Hz, 2H), 3.20 (s, 2H), 3.80 (brs, 1H), 4.37 (s, 2H), 4.86 – 4.90 (m, 2H), 6.50 (d, *J* = 8.1 Hz, 1H), 6.55 (d, *J* = 8.1 Hz, 1H), 7.05 (t, *J* = 8.1 Hz, 1H), 7.29 – 7.41 (m, 5H);

Molecular weight : 249.36

MS (M+H): 250

To a solution of N-benzyl-7-methylene-5,6,7,8-tetrahydronaphthalen-1-amine (0.50 g, 2.00 mmol) in tetrahydrofuran (5 ml) was added 0.5 M tetrahydrofuran solution of 9-borabicyclo[3.3.1]nonane dimer (8.20 ml, 4.10 mmol) at 0°C and then stirred at room temperature for 8 hours. To the resulting mixture was added 3N aqueous solution of sodium hydroxide (82 ml) followed by aqueous 33% hydrogen peroxide solution (2 ml), and the mixture was stirred at room temperature for 6 hours. The mixture was extracted with ethylacetate, and the organic layer was washed with brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (eluent: ethylacetate / hexane = 1 / 4) to provide N-benzyl-7-methyl-5,6,7,8-tetrahydronaphthalen-1-amine (0.069 g).

¹H NMR (CDCl₃): δ 1.11 (d, *J* = 6.9 Hz, 3H), 1.22 – 1.24 (m, 1H), 1.55 – 1.65 (m, 1H), 1.80 – 2.05 (m, 2H), 2.74 – 2.76 (m, 2H), 4.36 (s, 2H), 4.45 (brs, 1H), 5.50 (brs, 1H), 6.49 (m, 2H), 7.05 (t, *J* = 9.0 Hz, 1H), 7.28 – 7.40 (m, 5H).

A mixture of N-benzyl-7-methyl-5,6,7,8-tetrahydronaphthalen-1-amine (90.0 mg, 0.346 mmol) and palladium carbon (10.0 mg) in ethylacetate (10 ml) was stirred under hydrogen for 1 hour. The mixture was filtered through a pad of celite, and the filtrate was concentrated under reduced pressure. The obtained residue was purified by column chromatography (eluent: ethylacetate / hexane = 1 / 3) to provide (7-methyl-5,6,7,8-tetrahydronaphthalen-1-yl)amine (46.0 mg).


^1H NMR (CDCl_3): δ 1.05 (d, $J = 6.0$ Hz, 3H), 1.09 – 1.19 (m, 1H), 1.46 – 1.84 (m, 3H), 2.63 – 2.6 (m, 2H), 4.25 (brs, 2H), 4.36 (brs, 1H), 6.45 – 6.49 (m, 2H), 6.93 (t, $J = 6.0$ Hz, 1H);

Molecular weight : 161.25

MS ($\text{M}+\text{H}$): 162

5 Example 3-1

N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(7-methyl-5,6,7,8-tetrahydronaphthalen-1-yl)urea



A mixture of (7-methyl-5,6,7,8-tetrahydronaphthalen-1-yl)amine (30.0 mg, 0.186 mmol) and 4-chloro-3-trifluoromethyl isocyanate (50.0 mg, 0.220 mmol) in tetrahydrofuran (10 ml) was stirred at room temperature for 16 hours. After the mixture was concentrated under reduced pressure, the obtained residue was purified by silica gel column chromatography (eluent: ethylacetate / hexane = 1 / 3) to provide N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-(7-methyl-5,6,7,8-tetrahydronaphthalen-1-yl)urea (42.0 mg).

^1H NMR ($\text{MeOD}-d_3$): δ 1.10 (d, $J = 6.0$ Hz, 3H), 1.35 – 1.67 (m, 1H), 1.70 – 1.92 (m, 1H), 1.93 – 2.15 (m, 3H), 2.61 – 2.70 (m, 2H), 3.88 – 3.92 (m, 1H), 4.39 (d, $J = 6.0$ Hz, 1H), 6.86 (d, $J = 9.0$ Hz, 1H), 7.15 (d, $J = 9.0$ Hz, 1H), 7.48 (d, $J = 9.0$ Hz, 1H), 7.59 – 7.66 (m, 2H), 8.00 (s, 1H).


Molecular weight : 382.82

MS ($\text{M}+\text{H}$): 383

20 Activity Class : A

Starting material

(8-Amino-1,2,3,4-tetrahydronaphthalen-2-yl)methanol



To a solution of 8-amino-3,4-dihydronaphthalen-2(1H)-one (5.00 g, 31.0 mmol) and pyridine (3.68 g, 46.5 mmol) in tetrahydrofuran (60 ml) was added benzyl chloroformate (6.35 g, 37.2 mmol) at 0°C. After the mixture was stirred at room temperature for 1 hour, it was poured into water and extracted with ethylacetate. The organic layer was washed with brine, dried over MgSO₄, filtered, and concentrated under reduced pressure. The obtained residue was washed with diethylether to provide benzyl (7-oxo-5,6,7,8-tetrahydronaphthalen-1-yl)carbamate (6.52 g).

¹H NMR (CDCl₃): δ 2.58 (t, *J* = 6.8 Hz, 2H), 3.08 (t, *J* = 6.8 Hz, 2H), 3.47 (s, 2H), 5.19 (s, 2H), 6.37 (brs, 1H), 7.07 (d, *J* = 7.3 Hz, 1H), 7.22 (t, *J* = 7.9 Hz, 1H), 7.33 – 7.50 (m, 6H).

To 2.6 M solution of n-butyllithium in hexane (1.72 ml) cooled at 0 °C was added diisopropylamine (452 mg, 4.47 mmol) dropwise. After the mixture was stirred at room temperature for 15 minutes, a solution of benzyl (7-oxo-5,6,7,8-tetrahydronaphthalen-1-yl)carbamate (600 mg, 2.03 mmol) in tetrahydrofuran (1 ml) at –78°C and stirred for 1 hour. A solution of methoxymethyl(diphenyl)phosphine (550 mg, 2.23 mmol) in tetrahydrofuran (1 mL) was added to the reaction mixture at –78°C and then stirred for 16 hours at room temperature. The resulting mixture was poured into water and extracted with ethylacetate. The organic layer was washed with brine, dried over MgSO₄, filtered, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (eluent: hexane / ethylacetate = 10 / 1) to provide benzyl [(7E)-7-(methoxymethylene)-5,6,7,8-tetrahydronaphthalen-1-yl]carbamate (109 mg).


¹H NMR (CDCl₃): δ 2.25 (t, *J* = 6.3 Hz, 2H), 2.79 (t, *J* = 6.3 Hz, 2H), 3.33 (s, 2H), 3.60 (s, 3H), 5.21 (s, 2H), 5.95 (s, 1H), 6.49 (brs, 1H), 6.85 (d, *J* = 7.6 Hz, 1H), 7.12 (t, *J* = 7.9 Hz, 1H), 7.33 - 7.43 (m, 5H), 7.73 (brs, 1H).

5 A solution of benzyl [(7E)-7-(methoxymethylene)-5,6,7,8-tetrahydronaphthalen-1-yl]carbamate (51.0 mg, 0.16 mmol) in a mixture of tetrahydrofuran (3 ml) and 2N aqueous HCl (6 ml) was stirred at room temperature for 2 hours, and then extracted with ethylacetate. The organic layer was washed with brine, dried over MgSO₄, filtered, and concentrated under reduced pressure. The obtained residue was dissolved in ethanol (2 ml) and sodium borohydride (5.97 mg, 0.16 mmol) was added at room temperature. After stirred for 2 hours, the mixture was poured into water and
10 extracted with diethylether. The organic layer was washed with brine, dried over MgSO₄, filtered, and concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (eluent: hexane / ethylacetate = 4 / 1) to provide benzyl [7-(hydroxymethyl)-5,6,7,8-tetrahydronaphthalen-1-yl]carbamate (34.0 mg).

¹H NMR (CDCl₃): δ 1.39 (m, 1H), 1.48 (brs, 1H), 1.95 - 1.98 (m, 2H), 2.24 (dd, *J* = 16.1 Hz, 10.1 Hz, 1H), 2.72 (dd, *J* = 16.1 Hz, 5.2 Hz, 1H), 2.77 - 2.88 (m, 3H), 3.63 - 3.65 (m, 2H), 5.20 (s, 2H), 6.90 (d, *J* = 7.6 Hz, 1H), 7.13 (d, *J* = 7.7 Hz, 1H), 7.32 - 7.42 (m, 5H), 7.62 (brs, 1H).

A mixture of benzyl [7-(hydroxymethyl)-5,6,7,8-tetrahydronaphthalen-1-yl]carbamate (32.0 mg, 0.10 mmol) and palladium carbon (30 mg) in ethanol (2 ml) was stirred under hydrogen at room temperature for 16 hours. The resulting mixture was filtered through a pad of celite, and the filtrate
20 was concentrated under reduced pressure. The obtained residue was purified by silica gel column chromatography (eluent: hexane / ethylacetate = 4 / 1) to provide (8-amino-1,2,3,4-tetrahydronaphthalen-2-yl)methanol (11.0 mg).

¹H NMR (CDCl₃): δ 1.37 - 1.44 (m, 2H), 1.94 - 2.04 (m, 2H), 2.14 (dd, *J* = 16.1 Hz, 10.4 Hz, 1H), 2.64 (dd, *J* = 15.7 Hz, 15.3 Hz, 1H), 2.79 - 2.82 (m, 2H), 3.58 (brs, 2H), 3.69 (d, *J* = 15.3 Hz, 2H),
25 6.53 (d, *J* = 7.9 Hz, 1H), 6.57 (d, *J* = 7.5 Hz, 1H), 6.95 (t, *J* = 15.3 Hz, 1H).

Example 4-1**N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-[7-(hydroxymethyl)-5,6,7,8-tetrahydronaphthalen-1-yl]urea**

- 5 A mixture of (8-amino-1,2,3,4-tetrahydronaphthalen-2-yl)methanol (11.0 mg, 0.06 mmol) and 4-chloro-3-trifluoromethylphenyl isocyanate (13.7 mg, 0.06 mmol) in 1,4-dioxane (2 ml) was stirred for 2 hours at 50°C. The resulting mixture was concentrated under reduced pressure, and the obtained residue was washed with diisopropyl ether to provide N-[4-chloro-3-(trifluoromethyl)phenyl]-N'-[7-(hydroxymethyl)-5,6,7,8-tetrahydronaphthalen-1-yl]urea (14.0 mg).
- 10 ¹H NMR (DMSO-*d*₆): δ 1.31 (m, 1H), 1.75-1.83 (m, 1H), 1.85 - 1.91 (m, 1H), 2.21 (dd, *J* = 16.4 Hz, 10.4 Hz, 1H), 2.70-2.81 (m, 3H), 3.44 (t, *J* = 5.7 Hz, 1H), 4.67 (t, *J* = 5.1 Hz, 1H), 6.83 (d, *J* = 7.5 Hz, 1H), 7.05 (t, *J* = 7.8 Hz, 1H), 7.56 - 7.64 (m, 3H), 7.94 (s, 1H), 8.10 (d, *J* = 2.2 Hz, 1H), 9.49 (s, 1H).

mp 194 - 196°C;


- 15 Molecular weight : 398.81

MS (M+H): 399

Activity Class : A

Claims

1. An urea derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof:



5 wherein

n represents an integer of 0 to 6;

Q₁ and Q₄ independently represent direct bond or methylene;

Chemical bond between Q₂---Q₃ is selected from the group consisting of a single bond and a double bond;

10 when Q₂---Q₃ is a single bond, Q₂ represents CHR², or CO, and Q₃ represents CHR³,

when Q₂---Q₃ is a double bond, Q₂ represents CR² and Q₃ represents CR³;

wherein

R² represents hydrogen, hydroxy, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy;

15 R³ represents hydrogen, hydroxy, C₁₋₆ alkoxy, C₁₋₆ alkanoyloxy, or C₁₋₆ alkyl optionally substituted by hydroxy, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy,

with the proviso that Q₁ and Q₄ can not be direct bond at the same time;

R² and R³ can not be hydrogen at the same time;

when Q₁ and Q₄ are both methylene and R³ is hydroxy, R² is hydroxy, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy;

20 when Q₁ is direct bond, R² is hydroxy, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy; and when Q₄ is direct bond, R² is hydrogen, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy;

and

5 R^4 represents aryl optionally having one or two substituents selected from the group consisting of halogen, hydroxy, C_{1-6} alkylamino, $di(C_{1-6}$ alkyl)amino, C_{3-8} cycloalkylamino, C_{1-6} alkoxy carbonyl, phenyl, benzyl, sulfonamide, C_{1-6} alkanoyl, C_{1-6} alkanoylamino, carbamoyl, C_{1-6} alkylcarbamoyl, cyano, C_{1-6} alkyl optionally substituted by cyano, C_{1-6} alkoxy carbonyl, or mono-, di-, or tri-halogen, C_{1-6} alkoxy optionally substituted by mono-, di-, or tri- halogen, phenoxy optionally substituted by halogen or C_{1-6} alkyl, and C_{1-6} alkylthio optionally substituted by mono-, di-, or tri- halogen.

10 2. The urea derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1,

wherein

Q_1 and Q_4 represent methylene;

Q_2 — Q_3 is a single bond;

15 Q_2 represents CHR^2 , or CO ,

wherein

R^2 represents hydroxy, C_{1-6} alkoxy or C_{1-6} alkanoyloxy; and

Q_3 represents CHR^3 ,

wherein

20 R^3 represents hydrogen, hydroxy, C_{1-6} alkoxy or C_{1-6} alkanoyloxy.

3. The urea derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1,

wherein

Q_1 represents methylene;

25 Q_4 represents direct bond;

Q_2 — Q_3 is a single bond;

Q₂ represents CHR² or CO,

wherein

R² represents hydrogen, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy; and

Q₃ represents CHR³,

5 wherein

R³ represents hydrogen, hydroxy, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy,

with the proviso that R² and R³ can not be hydrogen at the same time.

4. The urea derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1,

10 wherein

Q₁ represents direct bond;

Q₄ represents methylene;

Q₂—Q₃ is a single bond;

Q₂ represents CHR² or CO,

15 wherein

R² represents hydroxy, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy;

Q₃ represents CHR³,

wherein

R³ represents hydrogen, hydroxy, C₁₋₆ alkoxy or C₁₋₆ alkanoyloxy.

20 5. The urea derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1,

wherein

Q₁ and Q₄ represent methylene;

$Q_2\text{---}Q_3$ is a double bond;

Q_2 represents CR^2 ,

wherein

R^2 represents C_{1-6} alkoxy or C_{1-6} alkanoyloxy; and

5 Q_3 represents CR^3 ,

wherein

R^3 represents hydrogen, C_{1-6} alkoxy or C_{1-6} alkanoyloxy.

6. The urea derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1,

10 wherein

Q_1 and Q_4 represent methylene;

$Q_2\text{---}Q_3$ is a single bond or a double bond;

when $Q_2\text{---}Q_3$ is a single bond, Q_2 represents CH_2 and Q_3 represents CHR^3 ,

and when $Q_2\text{---}Q_3$ is a double bond, Q_2 represents CH and Q_3 represents CR^3 ,

15 wherein

R^3 represents C_{1-6} alkyl optionally substituted by hydroxy.

7. The urea derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1,

wherein

20 n represents an integer of 0 to 1; and

R^4 represents phenyl optionally substituted with one or more substituents selected from the group consisting of chloro, bromo, fluoro, nitro, methoxy, trifluoromethyl and trifluoromethoxy.

8. The urea derivative of the formula (I), its tautomeric or stereoisomeric form, or a salt thereof as claimed in claim 1, wherein said urea derivative of the formula (I) is selected from the group consisting of:
- 5 N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(6,7-dihydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)urea;
- N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(7-hydroxy-6-methoxy-5,6,7,8-tetrahydronaphthalen-1-yl)urea
- N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(6-hydroxy-7-methoxy-5,6,7,8-tetrahydronaphthalen-1-yl)urea;
- 10 4-[[[4-Chloro-3-(trifluoromethyl)phenyl]amino]carbonyl]amino]-2,3-dihydro-1H-inden-2-yl acetate;
- 4-[[[4-(Trifluoromethyl)benzyl]amino]carbonyl]amino]-2,3-dihydro-1H-inden-2-yl acetate;
- N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(1-hydroxy-2,3-dihydro-1H-inden-4-yl)urea;
- 15 N-[4-Chloro-3-(trifluoromethyl)phenyl]-N'-(6-hydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)urea; and
- N-(6-Hydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)-N'-[4-(trifluoromethyl)benzyl]urea.
9. A medicament comprising urea derivative of the formula (I), its tautomeric or stereoisomeric form, or a physiologically acceptable salt thereof as claimed in claim 1 in as an
- 20 active ingredient.
10. The medicament as claimed in claim 9, further comprising one or more pharmaceutically acceptable excipients.
11. The medicament as claimed in claim 9, wherein said urea derivative of the formula (I), its tautomeric or stereoisomeric form, or a physiologically acceptable salt thereof is a VR1
- 25 antagonist.
12. The medicament as claimed in claim 9 for the treatment and/or prevention of an urological disorder or disease.

13. The medicament as claimed in claim 12, wherein said urological disorder or disease is urge urinary incontinence or overactive bladder.
14. The medicament as claimed in claim 9 for the treatment and/or prevention of pain.
15. The medicament as claimed in claim 14, wherein said pain is chronic pain, neuropathic pain, postoperative pain, or rheumatoid arthritic pain.
16. The medicament as claimed in claim 9 for the treatment and/or prevention of a disorder or disease related to pain.
17. The medicament as claimed in claim 16, wherein said disorder or disease related to pain is neuralgia, neuropathies, algesia, nerve injury, ischaemia, neurodegeneration, or stroke.
18. The medicament as claimed in claim 9 for the treatment and/or prevention of an inflammatory disorder or disease.
19. The medicament as claimed in claim 18, wherein said inflammatory disorder or disease is asthma or COPD.
20. Use of compounds according to claim 1 for manufacturing a medicament for the treatment and/or prevention of an urological disorder or disease.
21. Use of compounds according to claim 1 for manufacturing a medicament for the treatment and/or prevention of pain.
22. Use of compounds according to claim 1 for manufacturing a medicament for the treatment and/or prevention of an inflammatory disorder or disease.
23. Process for controlling an urological disorder or disease in humans and animals by administration of a VR1-antagonistically effective amount of at least one compound according to claim 1.
24. Process for controlling pain in humans and animals by administration of a VR1-antagonistically effective amount of at least one compound according to claim 1.
25. Process for controlling an inflammatory disorder or disease in humans and animals by administration of a VR1-antagonistically effective amount of at least one compound according to claim 1.

EPO - Munich
41
15. Okt. 2003

UREA DERIVATIVES

ABSTRACT

This invention relates to urea derivatives and salts thereof which are useful as active ingredients of pharmaceutical preparations.

The urea derivatives of the present invention have an excellent activity as VR1 antagonists and are useful for the prophylaxis and treatment of diseases associated with VR1 activity, in particular for the treatment of urinary incontinence, urge urinary incontinence, overactive bladder, chronic pain, neuropathic pain, postoperative pain, rheumatoid arthritic pain, neuralgia, neuropathies, algesia, nerve injury, ischaemia, neurodegeneration, stroke, inflammatory disorders, asthma and COPD.

PCT/EP2004/011008



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.